

KNA342/J INTERGARD 343 GREY PART A

4

10/28/14

1.

1.1. INTERGARD 343 GREY PART A
KNA342/J

1.2.

1.3.

626-6

(8-6)

1.4.

055-632-6286(),055 586 2310()

055 587 6276()

055 586 2310()

055 586 2310()

2.

2.1.

3; H226

/

2;H315

/

2;H319

1;H317

3;H412

2.2.

11 , 12



H315
H317
H319
H412

[]:

P210 / / /
P260 / /
P261 / /가 / / /
P262 , ,
P264
P272
P273
P280 / / /

[]:

P301+310 : /
P302+352 :
P303+361+353 () :
/
P305+351+338 가 : .가
P321 ().
P331
P333+313 /
P337 :
P362
P363
P370 :
P378 , , ,

[]:

P403+233 가

[]:

P501 ()

2.3. PBT (,) vPvB (,)

3.

/	%	GHS	
EPOXY RESIN CAS No: 0025036-25-3	30-40	/ 2;H319 / 2;H315, 1;H317	[1]
xylene CAS No: 0001330-20-7	10-20	3; H226 - 4;H312 - 4;H332 / 2;H315 / 2;H319 -1 ;H336 - 1;H372	[1][2]
Aluminium, alkyls CAS No: 0007429-90-5	2.5-5	Water react. 2;H261 H250	[1][2]

Titanium dioxide CAS No: 0013463-67-7	2.5-5		[1][2]
Solvent naphtha (petroleum), light aromatic CAS No: 0064742-95-6	1-2.5	1;H304	[1]
n-Butanol CAS No: 0000071-36-3	1-2.5	3; H226 - 4;H302 -1 ;H335 / 2;H315 / 1;H318 -1 ;H336	[1][2]
Ethylbenzene CAS No: 0000100-41-4	1-2.5	2;H225 - 4;H332 - 3;H373 1;H304 / 2;H315 / 2;H319 -1 ;H335	[1][2]
Propylene glycol mono methyl ether CAS No: 0000107-98-2	1-2.5	3; H226 -1 ;H336	[1][2]
polyamide dispersion CAS No: 0055349-01-4	<1	1;H317 - 4;H413	[1]
Methanol CAS No: 0000067-56-1	<1	2;H225 - 3;H331 - 3;H311 - 3;H301 -1 1;H370	[1][2]
	30-40	---	---

- 1)
- 2) 가
- 3) PBT vPvB
16

4.

4.1.

가

가

4.2. 가 /

4.3.

5. ,

5.1.

; , , , .
.

Note; 가 .

가 .

5.2.

, , , 가 . :

5.3.

가 .
가 .

6.

6.1.

가 가 가 , 가

6.2.

가 .

6.3.

.8 .

, , 가 . (13 .)

가 .

가 .

, 가 , 가 . ,

7.

7.1.

가 .

가 (LEL) (OEL)

가

가

가 (LEL) (OEL)

7.2. ()

, 8

가

가

61

가

1

7.3. Specific end use(s)

가

, 가

. 3

Hot surfaces, Sparks,

가

(60% ,)

8.

8.1.

(OEL)

(ACGIH)

(ACGIH)

ppm

mg/m³

ppm

mg/m³

Aluminium, alkyls

2

Ethylbenzene

125

545

100

435

Methanol

250

310

200

260

n-Butanol			C50	C150
Propylene glycol mono methyl ether	150	540	100	360
Talc				2
Titanium dioxide				10
xylene	150	655	100	434

(P) (Peak exposure limit)
(R)
(Sk)
(Sen)
(Cat 1)
(Cat 2) 가
(Cat 3)

DNEL/PNEC

8.2.

가

가 , (visor)

(overall)

가

가 가

가

pH / (°C) (°C) 65
29
(= 1)
(,) / : 1.4 (n-Butanol)
: 6.6 (xylene)
(Pa)

1.37

n- / (Log Kow)

9.2.

10.

10.1.

10.2.

.(Section 7)

10.3.

가 가

10.4.

.(7 .)

10.5.

10.6.

가

11.

(OEL)

	LD50, mg/kg	LD50, mg/kg	LD50, mg/L/4hr	/ LD50, mg/L/4hr
Aluminium, alkyls - (7429-90-5)				
EPOXY RESIN - (25036-25-3)				
Ethylbenzene - (100-41-4)	3,500.00,	15,433.00,	17.20,	
Methanol - (67-56-1)	5,628.00,	15,800.00,	85.00,	
n-Butanol - (71-36-3)	2,292.00,	3,430.00,		
polyamide dispersion - (55349-01-4)				
Propylene glycol mono methyl ether - (107-98-2)	5,000.00,	13,000.00,		
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	6,800.00,	3,400.00,		
Titanium dioxide - (13463-67-7)	10,000.00,	10,000.00,		6.82,
xylene - (1330-20-7)	4,299.00,	1,548.00,		20.00,

()		
()		
()		
/	2	
/	2	
	1	
(1)		
()		

12.

12.1.

Dangerous Preparations Directive 1999/45/EC

가
(3)

	96 hr LC50 mg/l	49 hr EC50 mg/l	ErC50 mg/l
EPOXY RESIN - (25036-25-3)			
xylene - (1330-20-7)	3.30, Oncorhynchus mykiss	8.50, Palaemonetes pugio	100.00 (72 hr), Chlorococcales
Aluminium, alkyls - (7429-90-5)	0.12, Oncorhynchus mykiss	3.50, Daphnia magna	
Titanium dioxide - (13463-67-7)	1,000.00, Fundulus heteroclitus	5.50, Daphnia magna	5.83 (72 hr), Pseudokirchneriella subcapitata
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	9.22, Oncorhynchus mykiss	6.14, Daphnia magna	19.00 (72 hr), Selenastrum capricornutum
n-Butanol - (71-36-3)	1,376.00, Pimephales promelas	1,328.00, Daphnia magna	500.00 (96 hr), Scenedesmus subspicatus
Ethylbenzene - (100-41-4)	4.20, Oncorhynchus mykiss	2.93, Daphnia magna	3.60 (96 hr), Pseudokirchneriella subcapitata
Propylene glycol mono methyl ether - (107-98-2)	1,000.00, Oncorhynchus mykiss	500.00, Daphnia magna	1,000.00 (96 hr), Selenastrum capricornutum
polyamide dispersion - (55349-01-4)			
Methanol - (67-56-1)	100.00, Pimephales promelas	10,000.00, Daphnia magna	16.912 (96 hr), Ulva pertusa

12.2.

가

12.3.

12.4.

12.5. , , 가

PBT (,) vPvB (,)

12.6.

13.

13.1.

가

14.

14.1.

1263

14.2.

14.3.

1263, , 3, III, 3[Y]

EmS

F-E,S-E

ICAO/IATA 3

14.4. III

14.5.

:

IMDG :

14.6. 가 가

14.7. MARPOL73/78 Annex II IBC Code .

15.

4 , 2 , III

MSDS 8 .

- Aluminium, alkyls (0007429-90-5)
- n-Butanol (0000071-36-3)
- Ethylbenzene (0000100-41-4)
- Talc (0014807-96-6)
- Titanium dioxide (0013463-67-7)

(CMR):

- Silica(quartz) (0014808-60-7)
- ethanol (0000064-17-5)
- Ethylbenzene (0000100-41-4)
- Titanium dioxide (0013463-67-7)

:

- Aluminium, alkyls (0007429-90-5)
- n-Butanol (0000071-36-3)
- Ethylbenzene (0000100-41-4)
- Titanium dioxide (0013463-67-7)
- xylene (0001330-20-7)

:

- Aluminium, alkyls (0007429-90-5)
- n-Butanol (0000071-36-3)
- Ethylbenzene (0000100-41-4)
- xylene (0001330-20-7)

가 :

()

:

()

:

()

:

()

Group I:

()

Group II:

Aluminium, alkyls (0007429-90-5)

Ethylbenzene (0000100-41-4)

xylene (0001330-20-7)

:

()

()

:

Talc (0014807-96-6)

16.

: 10/28/2014

: 4

: 10/21/2010

MSDS KOSHA, NITE, ESIS, NLM, SIDS, IPCS, NCIS

SDS

Section 3 Phrases

H225

H226

H250

H261 가

H302

H304

H312

H315

H317

H318

H319

H332

H335

H336

H372

H373

This SDS is valid for 5 years from the revised date on page 1.



Akzo Nobel

가